



(tek-ahr-tuhs)

THE MOST IMPORTANT INFORMATION TO KNOW ABOUT TECARTUS®

TECARTUS may cause side effects that are life-threatening and can lead to death. Call or see your healthcare provider or get emergency help right away if you get any of the following:

- Fever (100.4°F/38°C or higher)
- Difficulty breathing
- Chills or shaking chills
- Confusion
- Dizziness or lightheadedness
- Severe nausea, vomiting, or diarrhea
- Fast or irregular heartbeat
- Severe fatigue or weakness

It is important to tell your healthcare provider that you received TECARTUS and to show them your TECARTUS Patient Wallet Card. Your healthcare provider may give you other medicines to treat your side effects.

ABOUT TECARTUS

TECARTUS is a treatment for adults with mantle cell lymphoma or acute lymphoblastic leukemia. It is used following disease progression while on or after other treatment. TECARTUS is different than other cancer medicines because it is made from your own white blood cells, which have been modified to recognize and attack your lymphoma cells.

BEFORE GETTING TECARTUS, TELL YOUR HEALTHCARE PROVIDER ABOUT ALL YOUR MEDICAL PROBLEMS, INCLUDING IF YOU HAVE OR HAVE HAD:

- Neurologic problems (such as seizures, stroke, or memory loss)
- Lung or breathing problems
- Heart problems
- Liver problems
- Kidney problems
- A recent or active infection

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

HOW YOU WILL RECEIVE TECARTUS

- Since TECARTUS is made from your own white blood cells, your blood will be collected by a process called “leukapheresis” (loo-kah-fur-ee-sis), which will concentrate your white blood cells.
- Your blood cells will be sent to a manufacturing center to make your TECARTUS.
- Before you get TECARTUS, you will get 3 days of chemotherapy to prepare your body.

IMPORTANT FACTS

This is only a brief summary of important information about TECARTUS and does not replace talking to your healthcare provider about your condition and your treatment.

- When your TECARTUS is ready, your healthcare provider will give it to you through a catheter placed into your vein (intravenous infusion). The infusion usually takes less than 30 minutes.
- You will be monitored where you received your treatment daily for at least 7 days for patients with MCL and at least 14 days for patients with ALL after the infusion.
- You should plan to stay close to the location where you received your treatment for at least 4 weeks after getting TECARTUS. Your healthcare provider will help you with any side effects that may occur.
- You may be hospitalized for side effects. Your healthcare provider will discharge you if your side effects are under control and it is safe for you to leave the hospital.
- Your healthcare provider will want to do blood tests to follow your progress. It is important that you do have your blood tested. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

WHAT TO AVOID AFTER RECEIVING TECARTUS

- Do not drive, operate heavy machinery, or do other dangerous things for 8 weeks after you get TECARTUS because the treatment can cause sleepiness, confusion, weakness, and temporary memory and coordination problems.
- Do not donate blood, organs, tissues, or cells for transplantation.

THE POSSIBLE OR REASONABLY LIKELY SIDE EFFECTS OF TECARTUS

The most common side effects of TECARTUS include:

- Fever (100.4°F/38°C or higher)
- Low white blood cells (can occur with a fever)
- Low red blood cells
- Low blood pressure (dizziness or lightheadedness, headache, feeling tired, short of breath)
- Fast heartbeat
- Confusion
- Difficulty speaking or slurred speech
- Nausea
- Diarrhea

TECARTUS may increase your risk of getting cancers including certain types of blood cancers. Your healthcare provider should monitor you for this.

These are not all the possible side effects of TECARTUS. Call your healthcare provider about any side effects that concern you. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

GET MORE INFORMATION

- This is only a brief summary of important information about TECARTUS. Talk to your healthcare provider to learn more.
- Visit www.TECARTUS.com or call 1-844-454-KITE (5483).

